

# ***Joint SOGC/CSEP Clinical Practice Guideline: Exercise in Pregnancy and the Postpartum Period***

## **— Principal Authors —**

*Gregory A.L. Davies, MD, FRCSC, Kingston ON*

*Larry A. Wolfe, PhD, FACSM, Kingston ON*

*Michelle F. Mottola, PhD, London ON*

*Catherine MacKinnon, MD, FRCSC, Brantford ON*

## **SOGC Clinical Practice Obstetrics Committee**

Catherine MacKinnon (Chair), MD, FRCSC, Brantford ON

Marc-Yvon Arsenault, MD, FRCSC, Montreal QC

Elias Bartellas, MD, FRCSC, St. John's NL

Yvonne Cargill, MD, FRCSC, Ottawa ON

Tom Gleason, MD, FRCSC, Edmonton AB

Stuart Iglesias, MD, Gibsons BC

Michael Klein, MD, CCFP, FCFP, FAAP, FPS, Vancouver BC

Marie-Jocelyne Martel, MD, FRCSC, Saskatoon SK

Anne Roggensack, MD, Kingston ON

Kathi Wilson, BSc, RM, Ilderton ON

## **Canadian Society for Exercise Physiology Board of Directors**

Philip Gardiner (President), PhD, Montreal QC

Terry Graham, PhD, Guelph ON

Robert Haennel, PhD, Regina SK

Richard Hughson, PhD, Waterloo ON

Duncan MacDougall, PhD, Westport ON

John McDermott, PhD, North York ON

Robert Ross, PhD, Kingston ON

Peter Tiidus, PhD, Waterloo ON

François Trudeau, PhD, Trois Rivières QC

Larry Wolfe, PhD, Kingston ON

---

This guideline has been reviewed by the Clinical Practice Obstetrics Committee and approved by the Executive and Council of the Society of Obstetricians and Gynaecologists of Canada, and approved by the Board of Directors of the Canadian Society for Exercise Physiology.

## *Joint SOGC/CSEP Clinical Practice Guideline: Exercise in Pregnancy and the Postpartum Period*

Gregory A.L. Davies<sup>1</sup>, Larry A. Wolfe<sup>2</sup>, Michelle F. Mottola<sup>3</sup>,  
and Catherine MacKinnon<sup>4</sup>

---

### Catalogue Data

Davies, G.A.L.; Wolfe, L.A.; Mottola, M.F.; and MacKinnon, C. (2003). Joint SOGC/CSEP Clinical Practice Guideline: Exercise in pregnancy and the postpartum period. **Can. J. Appl. Physiol.** 28(3): 329-341. © 2003 Canadian Society for Exercise Physiology.

---

**Key words:** fetus, neonate, outcomes, aerobic, strength

**Mots clés:** fœtus, nouveau-né, résultat clinique, aérobie, force

### Abstract/Résumé

**Objective:** To design Canadian guidelines advising obstetric care providers of the maternal, fetal, and neonatal implications of aerobic and strength-conditioning exercises in pregnancy. **Outcomes:** Knowledge of the impact of exercise on maternal, fetal, and neonatal morbidity, and of the maternal measures of fitness. **Evidence:** MEDLINE search from 1966 to 2002 for English-language articles related to studies of maternal aerobic and strength conditioning in a previously sedentary population, maternal aerobic and strength conditioning in a previously active population, impact of aerobic and strength conditioning on early and late pregnancy outcomes, impact of aerobic and strength conditioning on neonatal outcomes, as well as for review articles and meta-analyses related to exercise in pregnancy. **Values:** The evidence collected was reviewed by the Society of Obstetricians and Gynaecologists of Canada (SOGC Clinical Practice Obstetrics Committee) with representation from the Canadian Society for Exercise Physiology, and quantified using the evaluation of evidence guidelines developed by the Canadian Task Force on the Periodic Health Exam.

---

<sup>1</sup>Department of Obstetrics and Gynaecology, Queen's University, Kingston, ON, K7L 3N6; <sup>2</sup>School of Physical and Health Education and Department of Physiology, Queen's University, Kingston, ON, K7L 3N6; <sup>3</sup>School of Kinesiology and Department of Anatomy and Cell Biology, University of Western Ontario, London, ON, N6A 3K7; <sup>4</sup>Avenue Medical Centre, 221 Brant Ave., Brantford, ON, N3T 3J2.

**Recommendations:**

1. All women without contraindications should be encouraged to participate in aerobic and strength-conditioning exercises as part of a healthy lifestyle during their pregnancy. (II-1,2B)
2. Reasonable goals of aerobic conditioning in pregnancy should be to maintain a good fitness level throughout pregnancy without trying to reach peak fitness or train for an athletic competition. (II-1,2C)
3. Women should choose activities that will minimize the risk of loss of balance and fetal trauma. (III-C)
4. Women should be advised that adverse pregnancy or neonatal outcomes are not increased for exercising women. (II-1,2B)
5. Initiation of pelvic floor exercises in the immediate postpartum period may reduce the risk of future urinary incontinence. (II-1C)
6. Women should be advised that moderate exercise during lactation does not affect the quantity or composition of breast milk or impact infant growth. (I-A)

**Validation:** This guideline has been approved by the SOGC Clinical Practice Obstetrics Committee, the Executive and Council of SOGC, and the Board of Directors of the Canadian Society for Exercise Physiology. **Sponsors:** This guideline has been jointly sponsored by the Society of Obstetricians and Gynaecologists of Canada and the Canadian Society for Exercise Physiology.

**But:** L'élaboration de directives canadiennes à l'intention des professionnels de la santé oeuvrant en obstétrique au sujet des effets des programmes d'exercices aérobies et de force chez les femmes enceintes, les fœtus, et les nouveau-nés. **Résultat clinique:** Connaissances des conséquences de l'exercice sur la morbidité maternelle, fœtale, néonatale, et sur la condition physique maternelle. **Études:** Recherche dans la base de données Medline (1966 à 2002) d'articles sur les effets des programmes d'entraînement aérobique et à la force chez des femmes enceintes initialement sédentaires, les effets des programmes d'entraînement aérobique et à la force chez des femmes enceintes initialement actives, les effets des programmes d'entraînement aérobique et à la force sur l'évolution de la grossesse, les effets des programmes d'entraînement aérobique et à la force sur les nouveau-nés. Revue de la littérature et méta-analyse des effets de l'exercice sur la grossesse. **Valeurs:** Le Comité de pratique clinique en obstétrique de la Société des obstétriciens et gynécologues du Canada (SOGC) et des représentants de la Société canadienne de physiologie de l'exercice (SCPE) ont révisé les données recueillies, puis les ont quantifiées en fonction des critères d'évaluation des faits élaborés par le Groupe d'étude canadien sur l'examen médical périodique.

**Recommandations:**

1. Toute femmes pour qui il n'y a pas de contra-indication devrait être encouragée à développer de saines habitudes de vie durant leur grossesse en incluant la pratique d'exercices aérobies et d'exercices de force. (II-1,2B)
2. Le but d'un programme d'exercices aérobies est de maintenir une bonne condition physique tout au long de la grossesse sans chercher à atteindre des niveaux élevés ni à s'entraîner en vue d'une compétition. (II-1,2C)
3. On recommande aux femmes enceintes de choisir des exercices aérobies qui réduisent au minimum les pertes d'équilibre et les traumatismes au fœtus. (III-C)

4. *Les femmes devraient être informées que l'exercice n'augmente pas les risques de conséquences néfastes pendant la grossesse ou pour les nouveaux nés. (II-1,2B)*
5. *Il est bénéfique de commencer les exercices du plancher pelvien au tout début de la période postpartum afin de réduire le risque d'incontinence urinaire. (II-1C)*
6. *Les femmes devraient être informées que des exercices modérés chez la mère allaitante ne modifient ni la quantité ni la composition du lait et n'affecte pas la croissance du nouveau-né. (I-A)*

**Validation:** *Cette directive a été approuvée par le Comité de pratique clinique en obstétrique de la SOGC, le Comité exécutif et le Conseil de la Société des obstétriciens et gynécologues du Canada et le conseil d'administration de la Société canadienne de physiologie de l'exercice.*

**Parrainage:** *L'élaboration de cette directive s'est effectuée sous le parrainage de la Société des obstétriciens et gynécologues du Canada et de la Société canadienne de physiologie de l'exercice.*

## Introduction

Canadians are encouraged to include exercise as part of a healthy lifestyle.<sup>1</sup> Many women enter pregnancy with regular aerobic and strength-conditioning activities already a part of their daily lives. Other women see pregnancy as an opportunity to modify their lifestyles to include more health-conscious activities.

The traditional medical advice has been for exercising women to reduce their habitual levels of exertion in pregnancy and for nonexercising women to refrain from initiating strenuous exercise programs.<sup>2,3</sup> This advice was based on concerns that exercise could affect early and late pregnancy outcomes by increasing core body temperature during embryogenesis, increasing the risk of congenital anomalies, and shifting oxygenated blood and energy substrates to maternal skeletal muscle away from the developing fetus, leading to disturbances in growth.<sup>2,3</sup>

Early studies focusing on hard physical work combined with undernutrition and forced exercise in laboratory animals tended to support these concerns.<sup>4,5</sup> Other concerns included the risk of maternal musculoskeletal injury due to changes in posture and centre of gravity or fetoplacental injury due to blunt trauma or stress effects from sudden motions.<sup>6</sup> Recent investigations, focusing on both aerobic and strength-conditioning exercise regimens in pregnancy, have shown no increase in early pregnancy loss, late pregnancy complications, abnormal fetal growth, or adverse neonatal outcomes, suggesting that previous recommendations have been overly conservative.<sup>7-16</sup>

Women and their care providers should consider the risks of not participating in exercise activities during pregnancy, including loss of muscular and cardiovascular fitness, excessive maternal weight gain, higher risk of gestational diabetes or pregnancy-induced hypertension, development of varicose veins and deep vein thrombosis, a higher incidence of physical complaints such as dyspnea or low back pain, and poor psychological adjustment to the physical changes of pregnancy.<sup>17</sup>

These guidelines have been designed to aid Canadian women and their care providers as they discuss the relative merits of aerobic and strength conditioning in pregnancy and the postpartum period. The quality of evidence reported in these guidelines has been described using the Evaluation of Evidence criteria outlined in the Report of the Canadian Task Force on the Periodic Health Exam<sup>18</sup> (see boxed

### Quality of Evidence Assessment<sup>18</sup>

The quality of evidence reported in these guidelines has been described using the Evaluation of Evidence criteria outlined in the Report of the Canadian Task Force on the Periodic Health Exam.

1. Evidence obtained from at least one properly randomized controlled trial.

II-1. Evidence from well-designed controlled trials without randomization.

II-2. Evidence from well-designed cohort (prospective or retrospective) or case-control studies, preferably from more than one centre or research group.

II-3. Evidence obtained from comparisons between times or places with or without the intervention. Dramatic results in uncontrolled experiments (such as the results of treatment with penicillin in the 1940s) could also be included in this category.

III. Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

list above). The guidelines have been jointly sponsored by the Society of Obstetricians and Gynaecologists of Canada (SOGC) and the Canadian Society for Exercise Physiology (CSEP).

### CLASSIFICATION OF RECOMMENDATIONS<sup>18</sup>

Recommendations included in these guidelines have been adapted from the ranking method described in the Classification of Recommendations found in the Canadian Task Force on the Periodic Health Exam.

- A. There is good evidence to support the recommendation that the condition be specifically considered in a periodic health examination.
- B. There is fair evidence to support the recommendation that the condition be specifically considered in a periodic health examination.
- C. There is poor evidence regarding the inclusion or exclusion of the condition in a periodic health examination, but recommendations may be made on other grounds.
- D. There is fair evidence to support the recommendation that the condition not be considered in a periodic health examination.
- E. There is good evidence to support the recommendation that the condition be excluded from consideration in a periodic health examination.

### Who Should Exercise in Pregnancy?

In uncomplicated pregnancies, women with or without a previously sedentary lifestyle should be encouraged to participate in aerobic and strength-conditioning exercises as part of a healthy lifestyle.<sup>7,12,14,19-22</sup> (II-1,2 B) Women with complicated pregnancies have been discouraged from participating in exercise activities for fear of impacting the underlying disorder or maternal or fetal outcomes.<sup>23</sup> The conditions listed in Table 1 represent exclusion criteria for subjects participating in research studies.<sup>19-22</sup> Evidence specifically detailing the risks of exercise in pregnancy for women with these conditions is not available (III-C). “Relative contra-indications” refers to conditions in which risks may exceed benefits of regular physical activity. The woman’s decision to be physically active or not should be made with qualified medical advice.

**Table 1** Contraindications to Exercise in Pregnancy

Absolute Contraindications	Relative Contraindications
<ul style="list-style-type: none"> <li>• Ruptured membranes</li> <li>• Preterm labour</li> <li>• Hypertensive disorders of pregnancy</li> <li>• Incompetent cervix</li> <li>• Growth restricted fetus</li> <li>• High order multiple gestation (≥ triplets)</li> <li>• Placenta previa after 28th week</li> <li>• Persistent 2nd or 3rd trimester bleeding</li> <li>• Uncontrolled Type I diabetes, thyroid disease, or other serious cardiovascular, respiratory, or systemic disorder</li> </ul>	<ul style="list-style-type: none"> <li>• Previous spontaneous abortion</li> <li>• Previous preterm birth</li> <li>• Mild/moderate cardiovascular disorder</li> <li>• Mild/moderate respiratory disorder</li> <li>• Anemia (Hb &lt;100 g/L)</li> <li>• Malnutrition or eating disorder</li> <li>• Twin pregnancy after 28th week</li> <li>• Other significant medical conditions</li> </ul>

*Note:* Reprinted and modified with permission from the Canadian Society for Exercise Physiology.<sup>23</sup>

The Physical Activity Readiness Medical Examination for Pregnancy (PARmed-X for Pregnancy) is a tool developed by the Canadian Society for Exercise Physiology and endorsed by the Society of Obstetricians and Gynaecologists of Canada and Health Canada (and available through CSEP’s web site <http://www.csep.ca/forms.asp>) for screening women interested in participating in physical activity during pregnancy.<sup>23</sup> The PARmed-X for Pregnancy includes a questionnaire for women to complete, to supply their obstetric care providers with pertinent medical history and a recent patient activity profile. It provides women with practical prescriptions for participating in aerobic and strength-conditioning activities and includes a tear-away medical clearance form that can be completed by the obstetric provider and presented for participation in organized prenatal fitness activities.

## RECOMMENDATION 1

All women without contraindications should be encouraged to participate in aerobic and strength-conditioning exercises as part of a healthy lifestyle during their pregnancy. (II-1,2B)

### When and How to Start an Exercise Program

Many women find that the best time to start an exercise program is in the second trimester, when the nausea, vomiting, and profound fatigue of the first trimester have passed and before the physical limitations of the third trimester begin. Concerns about the teratogenic effect of high core body temperatures in the early first trimester have not been demonstrated in studies of exercising women.<sup>24–28</sup> Women who have been exercising prior to pregnancy may continue their exercise regimens throughout pregnancy using the guidelines outlined below.<sup>7,10–12,14</sup> (II-1,2B)

When starting an aerobic exercise program, previously sedentary women should begin with 15 minutes of continuous exercise three times a week, increasing gradually to 30-minute sessions four times a week.<sup>19,20,29–31</sup> Episodic maximal exercise by pregnant women in a research setting appears to be safe for mother and fetus.<sup>32,33</sup> Reasonable goals of aerobic conditioning in pregnancy would be to maintain a good fitness level throughout pregnancy without trying to reach peak fitness or train for an athletic competition. (II-1,2 C) Elite athletes who continue to train during pregnancy require supervision by an obstetric care provider with knowledge of the impact of strenuous exercise on maternal and fetal outcomes. Women with special needs may require a referral to a physiotherapist, exercise physiologist, or sports medicine specialist to develop an appropriate exercise program.

## RECOMMENDATION 2

Reasonable goals of aerobic conditioning in pregnancy should be to maintain a good fitness level throughout pregnancy without trying to reach peak fitness or train for an athletic competition. (II-1,2C)

Women should choose aerobic activities that will minimize the risk of loss of balance and fetal trauma. Brisk walking, stationary cycling, cross-country skiing, swimming, or aquafit are aerobic exercises that cause less trauma to the joints and ligaments and less bouncing up and down of the centre of gravity than running or jogging.<sup>34</sup> It is suggested that a warm-up and cool-down period be included in any exercise regimen. (III-C)

## RECOMMENDATION 3

Women should choose activities that will minimize the risk of loss of balance and fetal trauma. (III-C)

There is less evidence on strength conditioning and weight training in pregnancy.<sup>10,35</sup> Some women may experience symptomatic hypotension from compression of the vena cava by the pregnant uterus and should modify these exercises to avoid the supine position after approximately 16 weeks' gestation.<sup>36</sup> The ability to perform abdominal strengthening exercises may be impeded by the development of diastasis recti and associated abdominal muscle weakness.<sup>37–39</sup> (II-2C, III-C)

Stretching and strength-training exercises such as yoga and Pilates have not been studied in a pregnant population.

### Exercise Intensity

There is an increase of 10 to 15 beats per minute in resting heart rate in pregnancy.<sup>40,41</sup> However, at maximal exercise levels, there is a blunted heart rate response as compared to the nonpregnant state.<sup>40,41</sup> Therefore, it is suggested that the use of conventional heart rate target zones be modified to account for this reduction in maximal heart rate reserve.<sup>23,36</sup> (III-C) A modified version of the conventional age-corrected heart rate target zone can be found in Table 2.<sup>23,36</sup>

Other measures of exercise intensity include the “talk test” and a visual rating of perceived exertion (see Borg’s Rating, next page). As the “talk test” implies, the woman is exercising at a comfortable intensity if she is able to maintain a conversation during exercise; she should reduce the exercise intensity if this is not possible. Exercising women can also use a visual scale to assess their exercise intensity.<sup>20</sup> A target rating of 12 to 14 on Borg’s scale of perceived exertion is suggested during pregnancy.<sup>23,36,42</sup> (III-C)

**Table 2 Modified Heart Rate Target Zones for Aerobic Exercise in Pregnancy**<sup>23,36</sup>

Maternal Age	Heart Rate Target Zone (beats/min)	Heart Rate Target Zone (beats/10 sec)
Less than 20	140–155	23–26
20–29	135–150	22–25
30–39	130–145	21–24
40 or greater	125–140	20–23

*Note:* Reprinted with permission from the Canadian Society for Exercise Physiology.

### Safety Precautions

In addition to exercise, other components of a healthy lifestyle in pregnancy include good nutrition and abstinence from smoking, alcohol, and illicit drugs.<sup>43,44</sup> Some sport activities carry significant risk in pregnancy and are considered contraindicated. Women should not scuba dive in pregnancy, as the fetus is not protected from decompression sickness and gas embolism.<sup>45</sup> Other activities such as horseback riding, downhill skiing, ice hockey, gymnastics, and cycling may also put the fetus at risk of harm due to the trauma or loss of balance but are not contraindicated. (III-C) Under normal circumstances and with appropriate hydration, moderate exercise at altitudes up to 1,800 to 2,500 m (6,000 to 8,250 ft) does not appear to significantly alter maternal or fetal well-being. However, women should be wary of hiking in a location where they might fall. For those women who do not live at higher altitudes, and who are planning on exercising at altitudes above



**BORG'S RATING OF PERCEIVED EXERTION**<sup>42</sup>

6	
7	very, very light
8	
9	somewhat light
10	
11	fairly light
12	
13	somewhat hard
14	
15	hard
16	
17	very hard
18	
19	very, very hard
20	

*Note:* A rating of 12–14 is appropriate for most pregnant women.

2,500 m, appropriate acclimatization is required.<sup>46,47</sup> (II-2B) Women should discuss their specific sport activities with their obstetric care provider to clarify the risks and make modifications if necessary. Women should stop exercising and seek medical attention if they experience any of the symptoms listed below. (III-C)

- Excessive shortness of breath
- Chest pain
- Presyncope
- Painful uterine contractions
- Leakage of amniotic fluid
- Vaginal bleeding

**Outcomes of Exercise in Pregnancy**

Most trials of exercising women in pregnancy lack randomization and a sample size large enough to assess differences in maternal or fetal outcomes.<sup>22,48</sup> This does not imply that there should be no limits to exercise in pregnancy, but rather that the trials to date have

not demonstrated large differences in pregnancy outcomes, such as early pregnancy loss, birth weight, and preterm delivery rate.<sup>7-12,14</sup> Studies of neonatal outcomes have similar limitations in size and design and do not show any increased risk for the offspring of exercising women.<sup>13,15</sup>

**RECOMMENDATION 4**

Women should be advised that adverse pregnancy or neonatal outcomes are not increased for exercising women. (II-1,2B)

**Exercise in the Postpartum Period**

Depending on the mode of delivery, most types of exercise can be continued or resumed in the postpartum period. With the added fatigue of delivery and newborn care, some women may need to reduce the intensity or length of their exercise sessions. Women who have had Caesarean delivery may slowly increase their aerobic and strength training, depending on their level of discomfort and other complicating factors such as anemia or wound infection. The 6-week postpartum evaluation is an opportunity for women and their obstetric care providers to discuss these issues. Initiation of pelvic floor exercises in the immediate postpartum period may reduce the risk of future urinary incontinence.<sup>49,50</sup>

**RECOMMENDATION 5**

Initiation of pelvic floor exercises in the immediate postpartum period may reduce the risk of future urinary incontinence. (II-1C)

## Exercise and Breastfeeding

Breastfeeding is the best method of providing optimal nutrition, immunologic-based protection, and emotional nurturing for the growth and development of infants.<sup>51</sup> Therefore, exercise frequency and intensity should not interfere with a mother's ability to breastfeed. Although exercise does not negatively affect milk production or composition,<sup>52-54</sup> lactic acid has been shown to be increased in the breast milk of women exercising at maximal intensity, but not in those exercising at moderate level.<sup>55-58</sup> There is controversy as to whether this short-term increase in lactic acid makes the breast milk less palatable to the nursing infant.<sup>55,56, 58-61</sup> Mothers who find that their baby does not feed as well right after exercising may consider feeding the baby right before exercising (which may also make the breasts more comfortable during exercise), postponing feeding until 1 hour after exercising, or expressing milk prior to exercising to be used after exercising. The growth of breastfeeding babies of exercising women is normal, even for the infants whose mothers are losing weight as part of their exercise regimen.<sup>53</sup>

### RECOMMENDATION 6

Women should be advised that moderate exercise during lactation does not affect the quantity or composition of breast milk or impact infant growth. (I-A)

## Resources for the Pregnant Woman and Her Obstetric Provider

Pregnant women interested in participating in aerobic and strength-conditioning exercises in pregnancy can be referred to the following publications:

- *Active Living During Pregnancy*<sup>36</sup>
- *Nutrition for a Healthy Pregnancy: National Guidelines for the Childbearing Years*<sup>43</sup>
- *Healthy Beginnings: Your Handbook for Pregnancy and Birth*<sup>62</sup>

## References

1. **Handbook for Physical Activity Guide to Healthy Active Living.** [Egalement disponible en français sous le titre *Cahier d'accompagnement du Guide d'activité physique canadien pour une vie active saine*]. (1998). Ottawa: Health Canada. Also available online at <<http://www.hc-sc.gc.gc/hppb/paguide/>>. Cited Feb. 5, 2003.
2. American College of Obstetricians and Gynecologists [ACOG]. (1985). **Exercise During Pregnancy and the Postnatal Period.** Washington, DC: ACOG.
3. Shangold MM. (1989). Exercise during pregnancy: Current state of the art. **Can. Fam. Physician** 35: 1675-1689.
4. Tafari N, Naeye RL, Gobeze A. (1980). Effects of maternal undernutrition and heavy physical work during pregnancy on birth weight. **Br. J. Obstet. Gynaecol.** 87: 222-226.
5. Terada M. (1974). Effect of physical activity before pregnancy on fetuses of mice exercised forcibly during pregnancy. **Teratology** 10: 141-144.

6. Clapp JF, Little KD. (1995). The interaction between regular exercise and selected aspects of women's health. **Am. J. Obstet. Gynecol.** 173: 2-9.
7. Clapp JF. (1989). The effects of maternal exercise on early pregnancy outcome. **Am. J. Obstet. Gynecol.** 161: 1453-1457.
8. Klebanoff MA, Shiono PH, Carey JC. (1990). The effect of physical activity during pregnancy on preterm delivery and birth weight. **Am. J. Obstet. Gynecol.** 163: 1450-1456.
9. Kulpa PJ, White BM, Visscher R. (1987). Aerobic exercise in pregnancy. **Am. J. Obstet. Gynecol.** 156: 1395-1403.
10. Hall DC, Kaufmann DA. (1987). Effects of aerobic and strength conditioning on pregnancy outcomes. **Am. J. Obstet. Gynecol.** 157: 1199-1203.
11. Hatch MC, Shu X, McLean DE, Levin B, Begg M, Reuss L, et al. (1993). Maternal exercise during pregnancy, physical fitness, and fetal growth. **Am. J. Epidemiol.** 137: 1105-1114.
12. Kardel KR, Kase T. (1998). Training in pregnant women: Effects on fetal development and birth. **Am. J. Obstet. Gynecol.** 178: 280-286.
13. Clapp JF, Lopez B, Harcar-Sevcik R. (1999). Neonatal behavioral profile of the offspring of women who continued to exercise regularly throughout pregnancy. **Am. J. Obstet. Gynecol.** 180: 91-94.
14. Sternfeld B, Queensberry CP, Eskenazi B, Newman LA. (1995). Exercise during pregnancy and pregnancy outcome. **Med. Sci. Sports Exerc.** 27: 634-640.
15. Clapp JF, Simonian S, Lopez B, Appleby-Wineberg S, Harcar-Sevcik R. (1998). The one-year morphometric and neurodevelopmental outcome of the offspring of women who continued to exercise regularly during pregnancy. **Am. J. Obstet. Gynecol.** 178: 594-599.
16. O'Neill ME. (1996). Maternal rectal temperature and fetal heart rate responses to upright cycling in late pregnancy. **Br. J. Sports Med.** 30: 32-35.
17. Wolfe LA, Mottola MF. (2000). Validation of guidelines for aerobic exercise in pregnancy. In: Kumbhare DA, Basmajian JV (Eds.), **Decision-Making and Outcomes in Sports Rehabilitation**, pp. 205-222. New York: Churchill Livingstone.
18. Wolf SH, Battista RN, Angerson GM, Logan AG, Eel W. (1994). **Canadian Task Force on the Periodic Health Exam**, p. xxxvii. Ottawa: Canada Communication Group.
19. Brenner IKM, Wolfe LA, Monga M, McGrath MJ. (1999). Physical conditioning effects on fetal heart rate responses to graded maternal exercise. **Med. Sci. Sports Exerc.** 31: 792-799.
20. Ohtake PJ, Wolfe LA. (1998). Physical conditioning attenuates respiratory responses to steady-state exercise in late gestation. **Med. Sci. Sports Exerc.** 30: 17-27.
21. Wolfe LA, Hall P, Webb KA, Goodman L, Monga M, McGrath MJ. (1989). Prescription of aerobic exercise during pregnancy. **Sports Med.** 8: 273-301.
22. Lokey EA, Tran ZV, Wells CL, Myers BC, Tran AC. (1991). Effects of physical exercise on pregnancy outcomes: A meta-analytic review. **Med. Sci. Sports Exerc.** 23: 1234-1239.
23. **Physical Activity Readiness Medical Examination for Pregnancy** [PARmed-X for pregnancy]. (2002). Ottawa: Canadian Society for Exercise Physiology. Available online at <<http://www.csep.ca/forms.asp>>. Cited May 2, 2003.
24. Kilham L, Ferm VH. (1976). Exencephaly in fetal hamsters following exposure to hyperthermia. **Teratology** 14: 323-326.

25. Smith DW, Clarren SK, Harvey MAS. (1978). Hyperthermia as a possible teratogenic agent. **J. Pediatr.** 92: 878-883.
26. Miller P, Smith DW, Shephard TH. (1978). Maternal hyperthermia as a possible cause of anencephaly. **Lancet** 1: 519-521.
27. Jones RL, Botti JJ, Anderson WM, Bennett NL. (1985). Thermoregulation during aerobic exercise in pregnancy. **Obstet Gynecol.** 65: 340-344.
28. Clapp JF. (1991). The changing thermal response to endurance exercise during pregnancy. **Am. J. Obstet. Gynecol.** 165: 1684-1689.
29. Webb KA, Wolfe LA, McGrath MJ. (1974). Effects of acute and chronic maternal exercise on fetal heart rate. **J. Appl. Physiol.** 77: 2207-2213.
30. Wolfe LA, Preston RJ, Burggraf GW, McGrath MJ. (1998). Effects of pregnancy and chronic exercise on maternal cardiac structure and function. **Can. J. Physiol. Pharm.** 77: 909-917.
31. Wolfe LA, Walker RMC, Bonen A, McGrath MJ. (1994). Effects of pregnancy and chronic exercise on respiratory responses to graded exercise. **J. Appl. Physiol.** 76: 1928-1936.
32. Heenan AP, Wolfe LA, Davies GAL. (2001). Maximal exercise testing in late gestation: Maternal responses. **Obstet. Gynecol.** 97: 127-143.
33. MacPhail A, Davies GAL, Victory R, Wolfe LA. (2000). Maximal exercise testing in late gestation: Fetal responses. **Obstet. Gynecol.** 96: 565-570.
34. Wolfe LA. (2000). Pregnant women and endurance exercise. In: Shephard RJ, Åstrand PO (Eds.), **Endurance in Sport**, 2nd ed., pp. 531-546. London: Blackwell Scientific.
35. Avery ND, Stocking KD, Tranmer JE, Davies GAL, Wolfe LA. (1999). Fetal responses to maternal strength conditioning exercises in late gestation. **Can. J. Appl. Physiol.** 24: 362-376.
36. Kochan-Vintinner A. (1999). Wolfe L, Mottola M. (Eds.), **Active Living During Pregnancy**. [Booklet. Includes advice on exercise techniques. Sponsored by Society of Obstetricians and Gynaecologists of Canada and Health Canada.] Ottawa: Canadian Society for Exercise Physiology. Also available online at <<http://www.csep.ca/publicationsmain.html>>. Cited Feb. 5, 2003.
37. Fast A, Weiss L, Ducommun EJ, Medina E, Butler JG. (1990). Low-back pain in pregnancy. Abdominal muscles, sit-up performance, and back pain. **Spine** 15: 28-30.
38. Boissonnault JS, Blaschak MJ. (1988). Incidence of diastasis recti abdominis during the childbearing year. **Phys. Ther.** 68: 1082-1086.
39. Gilleard WL, Brown JMM. (1996). Structure and function of the abdominal muscles in pregravid subjects during pregnancy and the immediate postbirth period. **Phys. Ther.** 76: 750-762.
40. Avery ND, Wolfe LA, Amara CE, Davies GAL, McGrath MJ. (2001). Effects of human pregnancy on cardiac autonomic function above and below the ventilatory threshold. **J. Appl. Physiol.** 90: 321-328.
41. Lotgering FK, Struijk PC, Van Doorne MB, Wallenburg HCS. (1992). Errors in predicting maximal oxygen consumption in pregnant women. **J. Appl. Physiol.** 72: 562-567.
42. Borg GAV. (1982). Psychophysical bases of perceived exertion. **Med. Sci. Sports Exerc.** 14: 377-381.
43. Health Canada. (1999). **Nutrition for a Healthy Pregnancy: National Guidelines for the Childbearing Years**. Sponsored by Society of Obstetricians and Gynaecologists

- of Canada, Dietitians of Canada, the College of Family Physicians of Canada, and the Federal/Provincial/Territorial Group on Nutrition. Ottawa: Minister of Public Works and Government Services Canada. Available online at <[http://www.hc-sc.gc.ca/hpfb-dgpsa/onpp-bppn/national\\_guidelines\\_int\\_e.html](http://www.hc-sc.gc.ca/hpfb-dgpsa/onpp-bppn/national_guidelines_int_e.html)>. Cited Feb. 5, 2003.
44. Gruslin-Giroux A, Selby P, Davies GAL, Leyland NA, Franche R. (1998). Alcohol abuse and the pregnant woman. **J. Soc. Obstet. Gynaecol. Can.** 20: 655-666.
  45. Camporesi EM. (1996). Diving and pregnancy. **Semin. Perinatol.** 20: 292-302.
  46. Artal R, Fortunato V, Welton A, Constantino N, Khodiguian N, Villabos L, et al. (1995). A comparison of cardiopulmonary adaptations to exercise in pregnancy at sea level and altitude. **Am. J. Obstet. Gynecol.** 175: 505-506.
  47. Huch R. (1996). Physical activity at altitude in pregnancy. **Semin. Perinatol.** 20: 303-314.
  48. Kramer MS (2001). Regular aerobic exercise during pregnancy [Cochrane review]. In: **The Cochrane Library**, 3. Oxford: Update Software.
  49. Morkved S, Bo K. (1996). The effect of post-natal exercises to strengthen the pelvic floor muscles. **Acta Obstet. Gynecol. Scand.** 75: 382-385.
  50. Morkved S, Bo K. (2000). Effect of postpartum pelvic floor muscle training in prevention and treatment of urinary incontinence: A one-year follow-up. **Br. J. Obstet. Gynaecol.** 107: 1022-1028.
  51. World Health Organization [WHO]. (1989). **Protecting, Promoting and Supporting Breastfeeding: The Special Role of Maternity Services**. Geneva: A Joint WHO/UNICEF Statement.
  52. Lovelady CA, Lonnerdal B, Dewey KG. (1990). Lactation performance of exercising women. **Am. J. Clin. Nutr.** 52: 103-109.
  53. Dewey KG, Lovelady CA, Nommsen-Rivers LA, McCrory MA, Lonnerdahl BA. (1994). A randomized study of the effects of aerobic exercise by lactating women on breast-milk volume and composition. **N. Engl. J. Med.** 330: 449-453.
  54. Prentice A. (1994). Should lactating women exercise? **Nutr. Rev.** 52: 358-360.
  55. Wallace JP, Inbar G, Ernsthansen K. (1992). Infant acceptance of post-exercise breast milk. **Pediatrics** 89: 1245-1247.
  56. Wallace JP, Rabin J. (1991). The concentration of lactic acid in breast milk following maximal exercise. **Int. J. Sports Med.** 12: 328-331.
  57. Quinn TJ, Carey GB. (1999). Does exercise intensity or diet influence lactic acid accumulation in breast milk? **Med. Sci. Sports Exerc.** 31: 105-110.
  58. Carey GB, Quinn TJ. (2001). Exercise and lactation. Are they compatible? **Can. J. Appl. Physiol.** 26: 55-74.
  59. Sampsel CM, Seng J, Yeo S, Killion C, Oakley D. (1999). Physical activity and postpartum well-being. **J. Obstet. Gynecol. Neonatal Nurs.** 28: 41-49.
  60. Duffy L. (1997). Breastfeeding after strenuous aerobic exercise: A case report. **J. Hum. Lact.** 13: 145-146.
  61. Wright KS, Quinn TJ, Carey GB. (2002). Infant acceptance of breast milk after maternal exercise. **Pediatrics** 109: 585-589.
  62. Schuurmans N, Lalonde A. (1998). **Healthy Beginnings: Your Handbook for Pregnancy and Birth**. [Antepartum, intrapartum, and postpartum issues including exercise.] Ottawa: Society of Obstetricians and Gynaecologists of Canada. Available online at <<http://www.healthy-beginnings.com/-healthybeginnings/>>. Cited Feb. 5, 2003.